

SPECTRUM OF THE MULTI-REGGEON COMPOUND STATES IN MULTI-COLOUR QCD *

Jan Kotański

M. Smoluchowski Institute of Physics, Jagellonian University
Reymonta 4, 30-059 Kraków, Poland

(Received June 30, 2002)

We study the properties of the colour-singlet compound states of reggeized gluons in multi-colour QCD. Applying the methods of integrable models, we calculate their spectrum and discuss the application of the obtained results to high-energy asymptotics of the scattering amplitudes in perturbative QCD.

The high-energy asymptotics of the scattering amplitude is governed by the t -channel exchange of arbitrary number $N = 2, 3, \dots$ of reggeized gluons described by the so-called “fish-net” or “checker-board” Feynman diagrams. The problem of resumming such diagrams formulated in sixties [1]. The lowest non-trivial contribution for $N = 2$ was calculated by Balitski, Fadin, Kuraev and Lipatov in 1978 [2], who derived and solved an equation for intercept of the compound state of $N = 2$ reggeized gluons – the BFKL Pomeron. The equation for three and more reggeized gluons was formulated in the 1980’s by Bartels, Kwieciński, Praszalowicz [3] and Jaroszewicz [4]. However, it took almost twenty years to obtain the solution for $N = 3$, which corresponds to the QCD odderon [5, 6]. Finally, the solutions for higher $N = 4, \dots, 8$ and their generalizations to arbitrary N were obtained recently in the multi-colour limit in series of papers [7, 8, 9] written in collaboration with S. É. Derkachov¹, G. P. Korchemsky², A. N. Manashov³.

The leading contribution to the total elastic scattering amplitude of two hadrons (A, B) can be written in QCD in the Regge limit: $s \rightarrow \infty$, $t = \text{const}$,

* Presented at the X International Workshop on Deep Inelastic Scattering (DIS2002)

¹ Department of Mathematics, St.-Petersburg Technology Institute, St.-Petersburg, Russia

² Laboratoire de Physique Théorique, Université de Paris XI, 91405 Orsay Cédex, France

³ Department d’ECM, Universitat de Barcelona, 08028 Barcelona, Spain

as a power series in a strong coupling constant $\bar{\alpha}_s = \alpha_s N_c / \pi$

$$\mathcal{A}(s, t) \sim -i \sum_{N=2}^{\infty} (i\bar{\alpha}_s)^N \frac{s^{1+\bar{\alpha}_s \epsilon_N}}{\sqrt{\bar{\alpha}_s \ln s}} \beta_A^{(N)}(t) \beta_B^{(N)}(t). \quad (1)$$

Above, N denotes the number of reggeized gluons, called Reggeons, propagating in the t -channel and the residue factors, $\beta_{A(B)}^{(N)}$, measure the overlap of the wave function of the compound state of N reggeized gluons with the wave functions of two scattered particles. The parameter ϵ_N is defined as the maximal energy for the Schrödinger (BKP) equation

$$\mathcal{H}_N \Psi(\{\vec{z}_k\}) = \epsilon_N \Psi(\{\vec{z}_k\}) \quad (2)$$

where $\Psi(\{\vec{z}_k\})$ is the Reggeon wave function and $\vec{z}_k$ denotes two-dimensional transverse coordinates of k^{th} reggeized gluon. $\mathcal{H}_N$ is the effective QCD Hamiltonian describing pair-wise interaction between N reggeons.

In the multi-colour limit, this Hamiltonian simplifies significantly [10, 11] leading to

$$\mathcal{H}_N = \sum_{k=0}^{N-1} H(\vec{z}_k, \vec{z}_{k+1}) \quad \text{where} \quad \vec{z}_0 \equiv \vec{z}_N. \quad (3)$$

It describes [8] the nearest neighbour interaction of the Reggeons and has a hidden cyclic and mirror permutational symmetry. Moreover, it possesses the set of the $(N-1)$ integrals of motion, which are the eigenvalues of conformal charges, $\hat{q}_k$ and $\hat{\bar{q}}_k$ ⁴ [7],

$$[\mathcal{H}_N, \hat{q}_n] = [\hat{q}_n, \hat{q}_m] = [\mathcal{H}_N, \hat{\bar{q}}_n] = [\hat{\bar{q}}_n, \hat{\bar{q}}_m] = 0, \quad n, m = 2, \dots, N. \quad (4)$$

Thus, this system is completely integrable. The lowest integral may be expressed in terms of the conformal $SL(2)$ weight of the state h as $q_2 = -h(h-1)$. The Hamiltonian (3) is equivalent to the XXX Heisenberg spin magnet [7].

Our solution of the Schrödinger equation (2) is based on the method of the Baxter Q-operator [7]. It relies on the existence of the operator $Q(u, \bar{u})$ depending on the pair of complex spectral parameters u and $\bar{u}$ and satisfying the following relations. It commutes with itself for different values of the spectral parameters and with the integrals of motion

$$[Q(u, \bar{u}), Q(v, \bar{v})] = [\hat{t}_N(u, \{\hat{q}_n\}), Q(v, \bar{v})] = [\hat{t}_N(\bar{u}, \{\hat{\bar{q}}_n\}), Q(v, \bar{v})] = 0, \quad (5)$$

⁴ *bar* does not denote complex conjugation for which we use an *asterisk*.

where

$$\hat{t}_N(u, \{\hat{q}_n\}) = 2u^N + \hat{q}_2 u^{N-1} + \dots + \hat{q}_N, \quad (6)$$

and u, v are two complex spectral parameters. It also has to satisfy the Baxter equations

$$\begin{aligned} \hat{t}_N(u, \{\hat{q}_n\})\mathbb{Q}(u, \bar{u}) &= u^N \mathbb{Q}(u+i, \bar{u}) + u^N \mathbb{Q}(u-i, \bar{u}) \\ \hat{t}_N(\bar{u}, \{\hat{q}_n\})\mathbb{Q}(u, \bar{u}) &= (\bar{u}+i)^N \mathbb{Q}(u, \bar{u}+i) + (\bar{u}-i)^N \mathbb{Q}(u, \bar{u}-i). \end{aligned} \quad (7)$$

Furthermore, the Q -operator has prescribed analytical properties, i.e. known pole structure, and asymptotic behaviour at infinity. The above conditions fix the Q -operator uniquely and allow us to quantize the integrals q_k [7, 13, 14, 15]. It turns out that it is possible to express the Hamiltonian (3) in terms of the Baxter Q -operator [7]. Combining together the solutions of the Baxter equations and the quantum conditions for q_k with the Schrödinger equation (2) we can calculate the energy spectrum.

For $N = 3$ there exist two integrals of motion, q_2 and q_3 . The quantized values of q_3 exhibit some structure that can be seen on Figure 1 (left panel) [9]. The circled crosses denote the ground states. They have the highest energy for fixed N . The spectrum of quantized q_3 at $N = 3$ may be




Fig. 1. Quantized values of the integrals of motion at $h = 1/2$ for different number of reggeons $N = 3$ (left) and $N = 4$ (right).

approximated by the following formula [12]

$$[q_3^{\text{approx}}(\ell_1, \ell_2)]^{1/3} = \frac{\Gamma^3(2/3)}{2\pi} \left(\frac{1}{2}\ell_1 + i\frac{\sqrt{3}}{2}\ell_2 \right) \quad (8)$$

where $\ell_1, \ell_2 \in \mathbb{Z}$ and $\ell_1 + \ell_2$ is even. Similar structure is observed for $N = 4$, $h = 1/2$ and $q_3 = 0$ (plot 1 on the right)

$$[q_4^{\text{approx}}(\ell_1, \ell_2)]^{1/4} = \frac{\Gamma^2(3/4)}{2\sqrt{\pi}} \left(\frac{1}{\sqrt{2}}\ell_1 + i \frac{1}{\sqrt{2}}\ell_2 \right). \quad (9)$$

All these states have continuation in $\nu_h = \text{Im}(h)$, so, the eigenvalues of $\{\hat{q}_k\}$ form linear trajectories in the full q_k space [16, 17]. The spectrum of q_N has similar structure for higher N .

N	iq_3	q_4	iq_5	q_6	iq_7	q_8	ϵ_N
2							2.7726
3	.20526						-.2472
4	0	.15359					.6742
5	.26768	.03945	.06024				-.1275
6	0	.28182	0	.07049			.3946
7	.31307	.07099	.12846	.00849	.01950		-.0814
8	0	.39117	0	.17908	0	.03043	.2810

Table 1. Quantum numbers q_k and energy, $\epsilon_N(\{q_k\})$, of the N -Reggeons states in the multi-colour QCD for $h = 1/2$.

For the ground states (tab. 1), the integrals of motion are either purely real or purely imaginary. Moreover, the odd integrals are equal to zero for even N 's. The numbers at $N = 2$ and $N = 3$ agree with the previously published ones [2, 5, 8, 15, 17], while the results for higher N are new. The energy is positive for even N 's and negative for odd N 's and it is shown in a plot 2. The contribution of these states to the scattering amplitude increases with s for the positive energy ϵ_N and decreases for the negative energy ϵ_N . The exact values of the energy are denoted by crosses on the left panel of fig. 2. The upper and the lower curves stand for the functions $1.8402/(N - 1.3143)$ and $-2.0594/(N - 1.0877)$, respectively.

The right panel in the figure 2 shows the dependence of the energy $\epsilon(\nu_h)$ along the ground state trajectory for different number of particles $2 \leq N \leq 8$. These functions are symmetrical in ν_h . The maximum energy is in $\nu_h = 0$. At large ν_h , $\epsilon_8 > \dots > \epsilon_3 > \epsilon_2$.

Summerizing we found the spectrum of the multi-Reggeon compound states in QCD. In the Pomeron sector (even N) the intercept of states is bigger than 1 but smaller than the intercept of the BFKL Pomeron: $\alpha_2 > \alpha_4 > \dots > 1$. In the odderon sector (odd N) the intercept of the states is smaller than 1 but it increases with N : $\alpha_3 < \alpha_5 < \dots < 1$.

Recently Lipatov and de Vega [18] found another set of the solutions for $N = 3, 4$ which differ from our expressions. This discrepancy requires further studies.




Fig. 2. The dependence of the ground state energy, ϵ_N , on the number of particles N and on the ν_h .

REFERENCES

- [1] S.C. Frautschi, *Regge Poles and S-matrix Theory*, New York, W.A. Benjamin, 1963; V.de Alfaro and T. Regge, *Potential Scattering*, Amsterdam, North-Holland, 1965; P.D.B. Collins, *An Introduction to the Regge Theory and High Energy Physics*, Cambridge University Press, 1977.
- [2] E.A. Kuraev, L.N. Lipatov and V.S. Fadin, *Phys. Lett.* **B60** (1975) 50; *Sov. Phys. JETP* **44** (1976) 443; **45** (1977) 199; I. Balitski, L.N. Lipatov, *Sov. J. Nucl. Phys.* **28** (1978) 822.
- [3] J. Bartels, *Nucl. Phys.* **B175** (1980) 365; J. Kwieciński and M. Praszalowicz, *Phys. Lett.* **B94** (1980) 413.
- [4] T. Jaroszewicz, High-energy multi-gluon exchange amplitudes, Trieste preprint IC/80/175, see also *Acta Phys. Pol.* **B11**, 965 (1980).
- [5] R. Janik and J. Wosiek, *Phys. Rev. Lett.* **79** (1997) 2935; *Phys. Rev. Lett.* **82** (1999) 1092.
- [6] L.Lukaszuk and B.Nicolescu, *Lett. Nuov. Cim.* **8** (1973) 405.
- [7] S. É. Derkachov, G. P. Korchemsky and A. N. Manashov, *Nucl. Phys.* **B617** (2001) 375
- [8] G. P. Korchemsky, J. Kotański, A. N. Manashov, *Phys. Rev. Lett.* **88** 122002.
- [9] S. É. Derkachov, G. P. Korchemsky, J. Kotański and A. N. Manashov [hep-th/0204124].
- [10] L.N. Lipatov, *JETP Lett.* **59** (1994) 596.
- [11] L.D. Faddeev, G.P. Korchemsky, *Phys. Lett.* **B342** (1995) 311; G.P. Korchemsky, *Nucl. Phys.* **B443** (1995) 255.
- [12] G. P. Korchemsky, *Nucl. Phys.* **B498** (1997) 68; preprint LPTHE-Orsay-97-73 [hep-ph/9801377].
- [13] Z. Maassarani and S. Wallon, *J. Phys.* **A28** (1995) 6423.
- [14] H.J. de Vega, L.N.Lipatov, *Phys. Rev.* **D64** (2001) 114019.

- [15] J. Kotański and M. Praszalowicz, *Acta. Phys. Pol.* **B33** (2002) 665.
- [16] M.A. Braun, St. Petersburg preprint SPbU-IP-1998/8 [hep-ph/9804432].
- [17] M. Praszalowicz and A. Rostworowski, *Acta Phys. Pol.* **B30**, 349 (1999).
- [18] H.J. de Vega, L.N. Lipatov [hep-ph/0204245].